

# TAMU COM Re-entry Student / Staff / Faculty

## POSITIVE COVID-19 Test<sup>1, 2</sup>

- \*Common COVID-19 Symptoms**
- Fever or chills
  - Cough
  - Shortness of breath
  - Fatigue
  - Muscle or body aches
  - Headache
  - New loss of taste or smell
  - Sore throat
  - Congestion or runny nose
  - Nausea or vomiting
  - Diarrhea

- Positive COVID-19 Test Result:**
- Exit in-person learning environment
  - Contact Supervisor
  - Complete COM COVID-19 Concern Form
  - Follow CDC Guidelines for self-quarantine
  - Must Log Temperature and Symptoms TWICE Daily
  - Complete TAMU COVID-19 Reporting Form

**Self-Isolation for  
MINIMUM of 10-Days  
(Day 1)**

**Never  
Symptomatic**

**COVID-19 Symptom Status  
DURING 10-Days of Self-Isolation**

**Symptomatic\*  
At Any Time  
(Start Isolation Over as Day 1)**

**Complete 10-Days  
Self-Isolation**

**Completed MINIMUM of 10-Days Self-Isolation  
Since Symptom Onset (Day 10 at Earliest)**

- No Fever > 100.3 F in the last 24 Hours
- Must be AFEBRILE WITHOUT Fever Reducing Medications
- Significant Improvement in Symptoms?

**Obtain Formal Medical  
Evaluation & Written  
Clearance from  
Personal PCP**

No ↓  
**Continue Self-Isolation  
Until Meeting Criteria  
Listed Above**

YES ↓  
**Obtain Formal Medical  
Evaluation & Written  
Clearance from Personal PCP**

Written Clearance  
Provided ↓

Written Clearance  
Provided ↓

**Contact Supervisor  
Regarding POSSIBLE  
Resumption of In  
Person Activities  
(Day 10)**

**Contact Supervisor Regarding POSSIBLE  
Resumption of In Person Activities  
(Day 10 at Earliest)**

**References -**

CDC guidelines for COVID positive Healthcare Providers (HCP) <https://www.cdc.gov/coronavirus/2019-ncov/hcp/return-to-work.html>

CDC guidelines for COVID positive and sick for EVERYONE <https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/isolation.htm>

CDC guidelines for COVID exposures for citizens and quarantine <https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/quarantine.html>

**FOOTNOTES -**

1 - Must adhere to CDC, TAMU, & hospital affiliate guidelines. More stringent local clinical affiliate guidelines supersede less restrictive COM guidelines & must be followed at all times.

2 – Guidance for ELEVATED-RISK / IMMUNOCOMPROMISED individuals may differ. Concurrent management with personal PCP is strongly advised.